

UCRL- 90425  
PREPRINT

MICROSTABILITY THEORY FOR DISTRIBUTIONS SEPARABLE  
IN ENERGY AND PITCH ANGLE

Gary R. Smith

This paper was prepared for submittal to  
the International Conference on Plasma  
Physics, Lausanne, Switzerland, June, 1984

March 20, 1984

Lawrence  
Livermore  
National  
Laboratory

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

## DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement recommendation, or favoring of the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

MICROSTABILITY THEORY FOR DISTRIBUTIONS SEPARABLE  
IN ENERGY AND PITCH ANGLE\*

Gary R. Smith

Lawrence Livermore National Laboratory

University of California

Livermore, California 94566, UNITED STATES OF AMERICA

**Abstract:** Distribution functions that are separable in energy and pitch angle allow analytical calculation of one or two velocity-space integrals that appear in the linear theory of certain microinstabilities.

### 1. Introduction

Linear stability theories of high-frequency waves in magnetized plasmas have usually relied on model particle distribution functions  $f$ . An oft-used model is

$$f = v_{\perp}^{2\ell} \exp(-\alpha_{\perp} v_{\perp}^2 - \alpha_{\parallel} v_{\parallel}^2) \quad , \quad \ell = 0, 1, 2, \dots \quad (1)$$

which is separable in the perpendicular and parallel velocity components  $v_{\perp}$  and  $v_{\parallel}$ :  $f = f_{\perp}(v_{\perp}) f_{\parallel}(v_{\parallel})$ .

(Normalization constants in  $f$  are omitted in this paper.)

This paper presents an additional class of model distributions that provide more realistic descriptions of some plasmas than does Eq. (1) or, more generally,  $f = f_{\perp} f_{\parallel}$ .

Model distributions allow one or two velocity-space integrations to be performed analytically, which greatly reduces the numerical work required to solve a dispersion relation. For example, for circularly polarized waves propagating along the ambient magnetic field ( $k_{\perp} = 0$ ) in a nonrelativistic plasma, one must calculate double velocity-space integrals like those in the susceptibility

$$\chi = \frac{\omega}{\omega - \Omega} + 2\pi \int_{-\infty}^{\infty} dv_{\parallel} \int_0^{\infty} dv_{\perp} v_{\perp} f \left[ \frac{k_{\parallel}^2 v_{\perp}^2}{2(\omega - \Omega - k_{\parallel} v_{\parallel})^2} + \left( \frac{\Omega}{\omega - \Omega} \right) \frac{k_{\parallel} v_{\parallel}}{\omega - \Omega - k_{\parallel} v_{\parallel}} \right] \quad . \quad (2)$$

For the distribution (1),

$$\chi = \frac{\omega}{\omega - \Omega} - \left[ \frac{(\ell + 1)\alpha_{\parallel}}{\alpha_1} + \frac{\Omega}{\omega - \Omega} \right] Z_1(\zeta) \quad , \quad (3)$$

$\zeta \equiv (\omega - \Omega)\alpha_{\parallel}^{1/2}/k_{\parallel}$ , and  $Z_1$  is the  $m = 1$  member of a family of functions ( $m = 0, 1, 2, \dots$ )

$$Z_m(\zeta) \equiv \pi^{-1/2} \int_{-\infty}^{\infty} du (u - \zeta)^{-1} u^m \exp(-u^2) \quad , \quad (4)$$

which are related to the standard plasma dispersion function  $Z_0$ . Numerical calculation of the right side of Eq. (3) involves evaluation of the analytic function  $Z_1$ , which is orders of magnitude quicker than the evaluation of the double integral in Eq. (2).

## 2. Nonrelativistic Plasmas with Symmetric Pitch-Angle Distributions

Consider now the class of distributions that are separable in speed  $v$  and pitch angle  $\phi = \cos^{-1}(v_{\parallel}/v)$ :  $f = F(v)G(\phi)$ . For nonrelativistic plasma and  $k_{\perp} = 0$ , the class of speed distributions

$$F(v) = v^{2\ell} \exp(-\alpha v^2) \quad , \quad \ell = 0, 1, 2, \dots \quad (5)$$

allows one of the two velocity-space integrals ( $\int_0^{\infty} dv v^2 \dots$ ) in  $\chi$  to be performed analytically. The pitch-angle integral can also be performed analytically if we choose

$$G = [H(\phi - \phi_+) - H(\phi - \phi_-) + H(\phi - \pi + \phi_-) - H(\phi - \pi + \phi_+)] \quad , \quad (6)$$

where  $H$  denotes the Heaviside step function. This  $G(\phi)$ , which obeys the symmetry  $G(\phi) = G(\pi - \phi)$ , is illustrated in Fig. 1. For  $\phi_- = \pi/2$ ,  $G(\phi)$  represents a distribution of




Fig. 1. The symmetric pitch-angle distribution given by Eq. (6).

particles with pitch angles in the range  $\phi_+ < \phi < \pi - \phi_+$  and models confinement in a magnetic mirror of ratio  $(\sin \phi_+)^{-2}$ . For  $\phi_- < \pi/2$ ,  $G(\phi)$  represents particles with pitch angles near  $\phi = \phi_i = (\phi_+ + \phi_-)/2$  and  $\phi = \pi - \phi_i$ , i.e., near a conical surface in velocity space. For the distribution given by Eqs. (5)-(6), the susceptibility (2) is

$$\chi = \frac{\omega}{\omega - \Omega} + \frac{\chi_+ \cos \phi_+ - \chi_- \cos \phi_-}{\cos \phi_+ - \cos \phi_-}$$

$$\chi_{\pm} = - \frac{1}{2 \left( \ell + \frac{1}{2} \right)!} \left[ \tan^2 \phi_{\pm} z_{2\ell+3}(\zeta_{\pm}) + \frac{\omega}{\omega - \Omega} \ell! \sum_{n=0}^{\ell} \frac{1}{n!} z_{2n+1}(\zeta_{\pm}) \right]$$

$$\zeta_{\pm} \equiv (\omega - \Omega) \alpha^{1/2} / k_{\parallel} \cos \phi_{\pm} \quad .$$

### 3. Nonrelativistic Plasmas with Unsymmetric Pitch-Angle Distributions

Pitch-angle distributions  $G(\phi)$  that are not symmetric about  $\phi = \pi/2$  but are piecewise constant in  $\phi$  also allow the pitch-angle integral to be performed analytically. An example is  $G(\phi) = H(\phi - \theta)$ ,  $0 \leq \phi \leq \pi$ , which represents a particle distribution that is isotropic except for the absence of all particles in an unsymmetric loss cone  $0 \leq \phi \leq \theta$ . For nonrelativistic plasma, for  $k_{\perp} = 0$ , and for the Maxwellian speed distribution (5) with  $\ell = 0$ , the susceptibility (2) is

$$\chi = \frac{\omega}{\omega - \Omega} \left( 1 - \frac{1}{(1 + \cos \theta)} \left\{ z_1(\zeta_0) - \bar{z}_1(\zeta_0) + [z_1(\zeta) + \bar{z}_1(\zeta)] \cos \theta \right\} - \frac{\sin \theta \tan \theta}{(1 + \cos \theta)} [z_3(\zeta) + \bar{z}_3(\zeta)] \right)$$

$$\zeta_0 \equiv (\omega - \Omega) \alpha^{1/2} / k_{\parallel} \quad , \quad \zeta \equiv \zeta_0 \sec \theta \quad .$$

The functions  $\bar{z}_1$  and  $\bar{z}_3$  are defined by ( $m = 0, 1, 2, \dots$ )

$$\bar{z}_m(\zeta) \equiv \pi^{-1/2} \int_{-\infty}^{\infty} du \operatorname{sgn}(u) (u - \zeta)^{-1} u^m \exp(-u^2) \quad ,$$

which differs from Eq. (4) only through the factor of  $\operatorname{sgn}(u)$ . Recursion relations exist among the  $\bar{z}_m$ , and

$$\bar{Z}_0(\zeta) = \pi^{-1/2} \exp(-\zeta^2) E_1(-\zeta^2) ,$$

where  $E_1$  is the exponential-integral function.

#### 4. Relativistic Plasmas

Plasmas with substantial fractions of relativistic electrons require the numerical calculation of one or two velocity-space integrals that appear in dispersion relations. This difficulty arises because of the complicated dependence on velocity-space variables of the wave-particle resonance conditions

$$\omega\gamma - n\Omega - k_{\parallel}p_{\parallel}/M = 0 , \quad (7)$$

where  $n$  is an integer,  $\gamma = [1 + (p_{\perp}^2 + p_{\parallel}^2)/M^2c^2]^{1/2}$ ,  $p_{\perp}$  and  $p_{\parallel}$  are momentum components, and  $M$  is the rest mass. One velocity-space integral (the pitch-angle integral) can be performed analytically if either  $k_{\perp} = 0$  or  $k_{\parallel} = 0$ , provided the distribution  $f$  has the form  $F(v)G(\phi)$  and  $G(\phi)$  is piecewise constant [e.g., Eq. (6)]. Since the speed integral is evaluated numerically, any  $F(v)$  can be used. Convenient velocity-space variables are  $x = \cos \phi$  and  $y = \gamma^{-1}$ , in terms of which Eq. (7) becomes

$$\omega - n\Omega y - k_{\parallel}cx (1 - y^2)^{1/2} = 0 . \quad (8)$$

The  $x$  dependence in Eq. (8) is simple enough to allow analytical calculation of the  $x$  (pitch-angle) integral if  $k_{\perp} = 0$ . For the other special wave-propagation direction ( $k_{\parallel} = 0$ ), Eq. (8) has no  $x$  dependence but the Bessel functions that appear for  $k_{\perp} \neq 0$  cause  $x$  dependence. Still, the  $x$  integral can be replaced by a rapidly convergent infinite sum and, in this sense, can be performed analytically.

---

\*Work performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under contract number W-7405-ENG-48.